

NTC Glossary

Tidal terminology

absolute sea level

When sea level is referenced to the centre of the Earth, it is sometimes referred to as “absolute”, as opposed to “relative”, which is referenced to a point (eg. a coastal benchmark, see § 5.9) whose vertical position may vary over time.

acoustic tide gauge *see* § 5.1

admittance *see* § 4.3

age of the tide

The delay in time between the transit of the moon and the highest spring tide. Normally one or two days, but it varies widely. In other words, in many places the maximum tidal range occurs one or two days after the new or full moon, and the minimum range occurs a day or two after first and third quarter. In a semi-diurnal tidal environment dominated by M_2 and S_2 , the age in hours can be computed using the formula $(g_{S_2} - g_{M_2})/(\omega_{M_2} - \omega_{S_2})$, where g_{S_2} and g_{M_2} are the **phase lags** (in degrees) from an analysis of the data, and ω_{M_2} and ω_{S_2} are the speeds (in °/hour). A similar formula can be devised for a diurnal environment dominated by O_1 and K_1 .

alias frequencies *see* § 5.3

amphidrome

Maps of specific tidal constituents (eg. M_2) are normally drawn with lines connecting points of constant amplitude (co-range or co-amplitude lines) and/or phase (co-phase lines) (Figure 6.1). The co-phase lines often appear like spokes radiating out from a central hub - the "amphidrome" or "node". Often a single map will show a number of these "amphidromic systems". The co-range lines more or less encircle the amphidrome, where the constituent amplitude is least. If M_2 is strongly dominant, the high water crest rotates around the amphidrome through the tidal cycle. The co-phase lines may be labelled in degrees or hours (eg 0 through 11 for S_2).




Figure 6.1 A schematic of an amphidrome. Dashed lines show amplitude, increasing away from the node. Solid lines show phase (degrees), increasing counter-clockwise around the amphidrome (typical of northern hemisphere amphidromes). If the co-phase lines were in hours instead of degrees, they would be labelled 0, 1, 2...11 for a semi-diurnal (approximately 30°/hour) tide, or 0, 2, 4...22 for a diurnal (approximately 15°/hour) tide.

Amphidromes are resonance phenomena, with higher-frequency constituents tending to have lesser areal extent. Whereas O_1 has an amphidrome virtually over the entire North Atlantic Ocean, M_2 has two.

amplitude *see* § 2.1

angular velocity

Rate of rotation, usually expressed in radians per unit time (as compared to the more familiar cycles per second). Since there are 2π radians per cycle, the angular velocity of Earth's rotation is 2π radians/sidereal day, or 0.729211×10^{-4} radians/second.

aphelion

The point in the elliptical orbit of Earth or other planet when it is furthest from the Sun.

apogee

The point in the Moon's elliptical orbit when it is furthest from earth. At this time, the tidal range tends to be reduced. The term "apogean" is sometimes used to indicate this situation, but its opposite, **perigean** seems more common.

apsides

The points in the orbit of a planet or moon which are the nearest and farthest from the centre of attraction. In the Earth's orbit these are called perihelion and aphelion, and in

the Moon's orbit, perigee and apogee. The line passing through the apsides of an orbit is called the line of apsides.

astronomical argument

The astronomical argument is essentially the same as the phase (§ 2.4), but omitting the term $i_a t$. For example, the solar day-based astronomical argument for the constituent M_1 is $-\lambda_s(t) + \lambda_h(t) + 90^\circ$. Based on lunar days, it is simply 90° . The corresponding astronomical arguments for N_2 are $-3\lambda_s(t) + 2\lambda_h(t) + \lambda_p(t)$ and $-\lambda_s(t) + \lambda_p(t)$. (In the shpNp' notation, which we recommend against, these are written $-3s + 2h + p$ and $-s + p$).

atmospheric tides *see* **radiational tides**

azimuth *see* **celestial sphere**

baroclinic/barotropic *see* **internal tides**

beat frequency *see* § 3.2

benchmark *see* § 5.8

black moon

A mythological object supposedly existing at the site of the unoccupied focus of the orbital ellipse traced out by the moon (the other focus being claimed by the earth). It has no other astronomical or tidal significance, but has a place in astrological mumbo jumbo.

bore *see* **tidal bore**

bubbler *see* § 5.1

cadastre

A register of land/marine boundaries. At the coastline, the cadastral boundaries are complicated by the changing nature of the waterline due to tides etc.

canal theory

An early mathematical attempt by Airy (1845) to explain the tidal motions by using the hydrodynamic equations of motion. The boundary conditions were unrealistic, treating the ocean as a canal with rigid vertical walls running around the equator or other latitude.

celestial sphere

Astronomers use the concept of a celestial sphere in order to have a reference system for locating objects in space. The points where the earth's axis of rotation intersects the celestial sphere are known as the *celestial poles*; the intersection of the plane containing the earth's equator is called the *celestial equator*. The angle between the celestial equator and a point on the sphere is the "declination" (north or south, as with latitude on earth). The angle along the celestial horizon measured between due north (or south if specified) from the observer clockwise to the point vertically below the point of interest is the "azimuth". The apparent path of the sun around the celestial sphere, over the course of earth's annual orbit, is known as the "ecliptic".




Figure 6.2 The celestial sphere, as seen from on high, far out in the starry heavens.

The green outer circle is a perimeter of the celestial sphere. The north and south celestial poles are labelled NCP and SCP. The plane of the ecliptic is yellow.

The ecliptic is generally used as a reference because most planets, our moon, and of course the sun all remain relatively close to the plane of the ecliptic, and hence their motions may be traced along or close to it (as opposed to the celestial equator, which is permanently inclined at 23.5°). This is why tables of orbital data for planets, for example, list the angle of inclination of their orbits to the *ecliptic*.

chart datum (CD) *see* § 5.9

circadian rhythms

Biological processes which re-occur on a regular basis governed by an internal timing mechanism are known as circadian rhythms. The timing may be reset by environmental changes such as changing length of day. Some animals also respond to tidal cycles, such as the spring-neap cycle. Some crabs have both circadian and tidal behavioural cycles, with their colour changing diurnally while their activity level varying over a period equal to the spring-neap cycle (even when removed to an aquarium). Related biological terms include circalunar (tied to the alignment of earth, moon, and sun) and circatidal (tied to the ebb and flood of the tide) rhythms, both pertaining to behaviour or physiology, which are usually found in littoral (nearshore oceanic) species.

component *see* § 5.8

compound tide *see* § 3.3

constants

Harmonic tidal analysis represents the sea level record as the sum of cosine waves. Each wave (or “constituent”) is uniquely identified by its frequency (or “speed”); for a given location, each frequency has an *amplitude* and *phase* (see § 2.1) which do not vary with time, and are hence known as “constants”. Tidal currents may also be harmonically represented by tidal constants, by first resolving them into north/south and east/west components (or along- and across-stream components).

constituent *see* constants

co-phase line and **co-range line** *see* amphidrome

Coriolis force

A moving body on the surface of the earth experiences a tendency to turn to the left (right) in the southern (northern) hemisphere due to earth's rotation. This tendency (which is an artifice of the rotating reference frame rather than an actual force) is known as the Coriolis force (or acceleration) is only noticeable with larger scale motions such as ocean currents and winds (despite the myth of bathtub drain vortices

rotating in opposite directions on either side of the equator). The Coriolis force affects the direction with which the tide propagates around an amphidrome and can also affect the propagation of the tide as it moves up a broad channel (most noticeably by tilting the water surface to the left or right of the direction of propagation).

co-tidal line

A line of constant phase or amplitude on a map of a tidal constituent (see **amphidrome**).

datum *see* § 5.8

day

The word day as commonly used refers to a mean solar day (msd) – the time between successive transits of the sun overhead or across a single meridian. The time required for earth to undergo a single revolution, known as a sidereal day, is slightly less owing to the earth’s orbital motion. A sidereal day is 0.9973 msd. The time between successive transits of the moon, known as the mean lunar day, is 1.035 msd - slightly longer than a msd as a consequence of the moon’s orbital motion.

declination *see* **celestial sphere**

diurnal tides *see* **species**

diurnal inequality

The condition whereby the daily high waters or low waters are of significantly different level.

dodge tide

Local South Australian term for a neap tide with minimal rise and fall over the course of a day or so. While very “flat” neaps (*see neap tide*) occur in a number of locations worldwide, the term “dodge” is used only in South Australia. Professor Sir Robert Chapman, C.M.G., writing in the Official Yearbook of the Commonwealth of Australia of 1938, stated “At spring tides the range, due to the semi-diurnal waves, is $2(M_2 + S_2)$, and at neaps, if the two are equal, or nearly equal, they practically neutralize one another and cause no rise nor fall at all. This is what happens at Port Adelaide where at this period the recording gauge shows frequently little or nothing in

the way of tide, in some cases the level of the water remaining almost constant for a whole day; in other cases one small tide occurs during the day. On each side of this tide is markedly irregular both as regards time and height, and the apparent impossibility of saying when the tide will be at this particular period has presumably gained for it its name 'The Dodger'."

The predicted times of high and low tides for Adelaide Outer Harbor on 10-11 August 2000 are given in Table 6.1.

Date	Time	Height (m)	Table 6.1 Instead of the normal
10	0304	1.2 (Low)	sequence of highs and lows, the tidal
10	1108	2.2 (High)	level remained virtually unchanged
10	2114	1.4 (Low)	from 2114 on 10 August until 0357 on
11	0248	1.5 (Dodge)	the following day. The next high tide
11	0357	1.5 (Dodge)	was not reached until nearly 11 hours
11	1415	2.2 (High)	later.
11	2158	1.1 (Low)	

This is put in context with the August, 2000 time series of sea level at Adelaide plotted below, with the dodge period encircled. Over most of the month the tidal behaviour is typical of semi-diurnal regimes worldwide. It is only the near-identity of the M_2 and S_2 amplitudes that give it its unique character at neaps.

The Canadian Department of Fisheries and Ocean website glossary defines a very similar phenomenon, which they call a "vanishing tide", defined thus: "the phenomenon occurring when a high and low water 'melt' together into a period of several hours with a nearly constant water level. The tide is in the diurnal category but is known as a 'vanishing tide'." An example of this may be found at Honiara, Solomon Islands, which exhibits a very flat period at neaps (Figure 6.3). Honiara has a diurnal regime (dominated by K_1 and O_1). During 5/6 August 2000, the predicted sea level variation remained within a 10 cm range for about nine hours, as opposed to a range at springs of about 90 cm. The moon entered its first quarter on 7 August.




Figure 6.3 Sea levels at Adelaide and Honiara. Intervals with virtually no tidal variation are encircled. In Adelaide these intervals are called the "Dodge Tide".

Doodson Numbers *see* § 2.3

earth tide

A deformation of the solid earth in response to the gravitational tidal forces of the sun and moon. The largest effect is the semi-diurnal deformation nearly in phase with the transit of the moon; its amplitude is less than 20 cm. Being nearly in phase with the tide-generating potential, and the absence of the resonance and dynamic features of ocean flow, mean that in some respects earth tides are closer to the **Equilibrium Tide** than are ocean tides. The solutions to the equations of motion for an elastic, spherical earth can be written in the form of vertical and horizontal displacements, plus a change in potential due to the deformation. Each of these is a simple linear function of the tide-generating potential, whose coefficients are known as the "Love numbers", h_2 (vertical displacement coefficient), l_2 (horizontal displacement coefficient), and k_2 (coefficient of change in the potential). Note that these refer to the deformation of the

solid earth with no ocean. A similar set of corrections to the ocean tides are described in **self-attraction and loading**.

ebb *see* **streams**

ecliptic *see* **celestial sphere**

epoch

The time origin used to reference the longitudes of astronomical features such as the lunar perigee. The word epoch is also used in at least two other very different ways in tidal work – as a synonym for **phase lag**, and for a period of time (usually a nodal cycle) over which a mean is calculated as the basis for a tidal datum.

equilibrium tide *see* § 1.3

equilibrium phase *see* § 1.3

eustatic sea level change

Global changes of sea level taking place over many years. Some authors associate "eustatic" with ocean volume changes, others with globally synchronous changes. However, such ocean surface displacements are now understood to be spatially irregular (in some areas, even opposing the global trend).

evection and variation

Two of many perturbations to the moon's orbit caused by changes in the solar gravitational potential during the course of the orbit, giving rise to the evectional (ρ_1 , X_1 , θ_1 , ν_2 and λ_2) and variational (σ_1 , μ_2) constituents.

establishment of a port

Definitions vary, but it is essentially the same as the more modern term, lunitidal interval.

extended harmonic method

This term usually refers to an tidal analysis of 114 or more terms. Prior to the work of Zetler and Cummings (1967) and Rossiter and Lennon (1968), tidal harmonic

analyses generally contained 64 or less constituents. Using spectral analysis, these authors independently identified an additional 54 constituents (individual terms differed between the two) which subsequently became a standard part of tidal analyses. The authors found that the reduction in the variance of the residuals following the inclusion of the additional terms was less than 10%.

flood *see* **streams**

form factor

A factor used to characterise the tides in an area as being predominantly diurnal, semi-diurnal, or mixed. Usually computed as $(H_{K1} + H_{O1}) / (H_{M2} + H_{S2})$, where H is the amplitude of the constituent in the subscript. The cut-off points are usually: less than 0.25, semidiurnal; 0.25 to 3.0, mixed; greater than 3.0, diurnal.

fortnightly tides

In most parts of the world, the tides go through a fortnightly "spring-neap" cycle, as described in Section 3.2. These are beat phenomena rather than actual harmonics. There are, however, harmonics arising directly from the **tide-generating potential** which have a period of a fortnight (two weeks), the most important being M_f . The harmonic MS_f arises from interactions occurring in shallow water (see § 3.3).

geoid *see* **geopotential**

geopotential

A gravitational field can be characterised by a "potential", the negative gradient of which defines the strength and direction of the force exerted upon a mass within the field. The earth's gravitational potential is called the geopotential. A geopotential surface is one whose potential is everywhere equal. In the absence of planetary rotation and forces other than earth's own gravity, the ocean would be at rest and its surface would conform to a geopotential surface known as the "geoid". Such forces include wind stress, density variations, and large-scale ocean waves. These may cause the mean sea level to differ (locally but semi-permanently) from the geoid by as much as a metre. The well-known 20 cm "head" of sea level between the Gulf of Panama and the Caribbean, caused by the difference in water densities, essentially means that

the geoid passing through mean sea level on the Caribbean side passes 20 cm below mean sea level on the Pacific side.

harmonic constants *see constants*

high water full and change (HWF&C)

Despite the name, HWF&C refers to a time interval, not a tidal plane. It is a somewhat antique term essentially synonymous with **lunitidal interval**. The “full and change” refers to full and new moon - the only part of the lunar cycle when it was useful. As with lunitidal interval, its purpose was to indicate the approximate delay following noon or midnight of the next high tide.

higher high water (HHW), highest astronomical tide (HAT), etc: *see § 5.9*

Indian spring high/low water *see § 5.9*

hydrodynamic equations of tidal motion *see § 1.4*

inference of constituents *see § 5.5*

internal tide

The ocean usually has a less dense upper layer overlying the much deeper, denser waters. Waves known as “internal waves” often occur on the interface between two such layers. If the interface is gradual, the direction of wave propagation may possess a vertical component, trapped by refraction within upper and lower limits. Internal waves are usually caused by flow in the lower layer moving over an obstacle such as an undersea ridge, with semi-diurnal tidal flows into and out of a fjord being a typical example. When this happens, a semi-diurnal internal wave, or “internal tide” is produced. Although these waves do not significantly affect the sea surface, they may be detected by satellite as bands of surface slicks due to the convergence of surface currents that are produced (Figure 6.4).



Figure 6.4 A NASA satellite photo showing internal waves. According to the website, "In the Sulu Sea between the Philippines and Malaysia, sunglint highlights delicate curving lines of internal waves moving to the northeast toward Palawan Island." Photo credits: NASA.

http://earthobservatory.nasa.gov/Newsroom/NewImages/images.php3?img_id=15334

intertidal zone

The part on a beach that lies between high and low tidal levels – sometimes exposed, and sometimes inundated, depending on the tide. For legal purposes, the high and low waters may be given more precise definitions, such as “mean high water” and “mean low water”.

inverse barometer effect *see* § 5.2

kappa phase *see* § 2.1

king tide

Term used colloquially in some parts of Australia and elsewhere for a seasonal high tide often combined with onshore winds, or any exceptionally high tide, in some cases due to a storm surge (see **storm surge**).

lagging of the tide *see* **lunitidal interval**

Loading Love numbers *see* **self-attraction and loading**

Love numbers *see* **earth tide**

low water (LW), lower low water (LLW), lowest astronomical tide (LAT), etc.:
see **tidal planes**

lunisolar tide

A tidal constituent whose origins are a combination of lunar and solar – that is, by coincidence, there being identical forcing frequencies stemming from both sources. The most important lunisolar tides are K_1 and K_2 . A shallow-water tide, M_{sf} , arises from the interaction between M_2 and S_2 .

lunitidal interval

The time interval between the moon's transit (overhead or below) and the following high tide. On the day of new or full moon, the moon's transit coincides with the sun's, providing a simple way to estimate the lunitidal interval – it is the number of hours after noon of the next high tide on that day. Formulae are sometimes given – for example, if the local phase lag of the moon's primary constituent, M_2 , is known, multiply it by 0.0345 (0.0345 being equal to the period of M_2 , 12.42 hours, divided by 360°). At Auckland, the phase lag of M_2 is 204° so the interval is seven hours according to the formula. A look at a tide table for Auckland reveals that at new and full moons, the first high tide after noon is, indeed, usually at about 7 p.m. At Outer Harbor, Adelaide, the M_2 phase is 106.6° , so according to the formula, the lunitidal interval is 3.7 hours. The formula (and in fact the concept of lunitidal interval) is of little practical value for locations where M_2 is not the dominant constituent. At Adelaide, where M_2 and S_2 share equal pre-eminence, the actual high tide at new and full moons comes about $4\frac{1}{2}$ to $5\frac{1}{2}$ hours after noon (and midnight) – essentially an average between the lunitidal and “solar-tidal” intervals (the latter being 6 hours).

Note: for reasons inscrutable to the practical man, some authors define lunitidal interval in terms of the prime meridian and local high water.

Since the lunar day is about 50 minutes longer than a solar day, the lunar wave arrives about 50 minutes later each day at a typical semi-diurnal port. On either side of a spring tide, this implies that the time of high tide first catches up with and then passes the lunitidal interval. Mariners (naturally) had a name for these decreasing and then increasing time delays: the "lagging" and "priming" of the tides, respectively.

mean high water (MHW), mean sea level (MSL), etc.: *see* **tidal planes**

meteorological tides *see* **radiational tides**

mixed tide *see* **form factor**

month

There are four types of month used in astronomy and relevant to the tidal gravitational potential. The moon completes a single orbit of the earth in a sidereal month, equal to 27.3217 days (mean solar days). During this time, the perigee has moved about 3° in its rotation of earth; consequently the time between the moon successively being at perigee is 27.5546 months (the *anomalistic month*). Similarly, during this time the lunar ascending node will have undergone a regression of about 4.5° (thereby reducing the length of time between successive passages of the moon through the ecliptic), thus defining the slightly shorter *nodical month* of 27.2122 days. A *synodic month* is the time between successive full moons. Because the earth progresses in its orbit around the sun while the moon is orbiting the earth, it takes longer than a sidereal month for the lunar phases to repeat. The synodic month is 29.5307 days.

NGWLMS *see* § 5.1

nodal cycle *see* § 3.4

nodal factor/phase *see* § 3.4

nonlinear tides *see* § 3.3

orthotide *see* § 4.3

overtide *see* § 3.3

perigeal cycle *see* **perigee**

perigean tide

Tides of increased range occurring monthly as the result of the Moon being in perigee. In some places (notably the Bay of Fundy) this modulation may equal that of the spring-neap cycle. The moon is at perigee every 27.5546 days, but the time between full moons is 29.5307 days. Thus, these two “beat” in and out of phase every 412 days. Since there are two spring tides per period between full moons, the perigean tide and the spring tides come into phase every 206 days. In terms of tidal analysis, the largest constituent due to the ellipticity of the lunar orbit is N_2 . There are also constituents associated with the ellipticity of the solar orbit, but because it is more circular than the lunar orbit (i.e., its eccentricity is less than a third), these constituents are much smaller. The opposite situation is known as apogean.

perigee

In the moon’s elliptical orbit around the earth, its point of closest approach is known as perigee. Over time, the orientation of the orbit *within* the orbital plane gradually rotates. As a consequence, the perigee circles the earth every 8.85 years, a period known as known as the perigeal cycle (not to be confused with perigean tide), and designated “p” in tidal literature. This is distinct from the nodal cycle, in which the orbital plane itself rotates. The moon is at perigee every 27.5546 days.

perihelion

The point of closest approach in the earth’s orbit around the sun. The sun is at perihelion every 365.2596 days – currently this nearly coincides with the start of the year (coincidentally midsummer in the southern hemisphere). The perihelion itself circles the sun every 20,942 years, in a rotation analogous to the perigeal cycle. The period is often designated "p' " or "p₁" in tidal literature.



Figure 6.5 Funafuti, Tuvalu – a photo of the southernmost islands of the atoll, viewed from the southeast. The nation of Tuvalu is comprised of nine coral atolls, the highest reaching an altitude of five metres. Tide gauges operating in Tuvalu (Figure 6.5) since the late 1970's have recorded a moderate rise in sea level (less than 3 cm), but **spring tides** inundate low-lying areas in the early part of most years when the earth is at **perihelion**. Photo credit: Allan Suskin.

phase lag *see* § 2.4

pole tide

A small tide of varying period (approximately 433 days, but varying) associated with changes in the earth axis of rotation known as the “Chandler Wobble”. Ultimately, it can be said to fall in the class of radiational tides, since the precession has been shown to be caused by oceanographic and meteorological variations, which redistribute water masses. This precession is independent of, and much smaller than, the precession of the equinoxes, which has a period of 26,000 years (see **year**). The largest reported pole tide is 30 mm, from the Gulf of Bothnia.

primary port *see* **standard port**

prime meridian

The meridian of 0° longitude, known also as the Greenwich Meridian.

priming of the tide *see* **lunitidal interval**

quadrature

The condition whereby the angle formed by the sun, earth, and moon is 90° . *See also* **syzygy**.

radiation stress

radiational tides

A quasi-periodic rise and fall of sea level caused by meteorological variability, hence also known as “meteorological tides”. Semi-diurnal radiational tides in the tropics are thought to be due to semi-diurnal fluctuations in surface barometric pressure forced at diurnal period at the top of the atmosphere (sometimes called "atmospheric tide"). Diurnal radiational tides are often caused by land/sea breezes or solar heating (note that neither of these forcing functions are purely sinusoidal in time). Monsoonal winds may cause semi-annual radiational tides on some coastlines. Annual heating of the atmosphere and redistribution of air mass can both cause annual radiational tides.

range

The difference between the maximum and minimum water levels during a typical tidal cycle.

Rayleigh criterion *see* § 5.4

rectilinear currents *see* **streams**

red tide

A discolouration of lake or sea water caused by an algal bloom having very little to do with tides.

regression of lunar nodes

Since the clockwise or western rotational direction of the lunar nodes around the **ecliptic** is opposite to that of most other rotations and orbits of the solar system, it said to be in regression.

relative sea level *see* **absolute sea level**

residuals *see* § 5.2

resonance

For any gulf or other body of water, there are certain resonant frequencies. These depend primarily on its dimensions (breadth and depth). If forced at the resonant frequency, water motions are amplified. For example, Spencer Gulf in South Australia is "tuned" to K_1 . When the K_1 wave enters the Gulf, its amplitude is about 3.3 times larger than P_1 (for which the Gulf is less well-tuned). The ratio increases up the Gulf, going from 3.3 to about 5.0 at the head (top end). The "ability" of Spencer Gulf to discriminate between two close frequencies (their periods differ by less than 8 minutes) attests to the fact that ocean systems are relatively lightly damped. Perhaps the most famous such case is the Bay of Fundy, also mentioned in the context of perigean tides.

response method *see* § 5.9

revolution without rotation

This intriguing phrase represents a useful simplification which is used in some elementary texts on tides, in their discussion of the balance of gravitational and centrifugal forces. According to that approach, the effect of the earth's daily rotation is ignored, leaving only the centrifugal force associated with "revolution", i.e. the earth-sun or earth-moon orbit. Actually, the term "ignored" is not completely correct. The diurnal rotation is accounted for in the gravitational field at the earth's surface (a vector field), which governs mean sea level.




Figure 6.6 Five successive views of the earth from above the North Pole.

Note that the path traced out by a point on the earth, symbolised by a small

black circle, never closes on itself. In fact, the waviness is exaggerated - the actual path would appear nearly straight.

If an observer in space above the North Pole watched an illuminated point on the surface of the earth (say, for example, at Kuala Lumpur), the point would trace a wavy line as in Figure 6.6 – not a series of closed loops as one might expect. This is because the earth travels through a distance of about 201 earth diameters every day on its orbit around the sun. For illustration, Figure 6 is drawn as if the distance were only eight earth diameters, so the true path would appear far less wavy than shown.

rip

A narrow shearing current flowing offshore through the surf zone. Sometimes (misleadingly) called a rip tide. Rips are a part of a circulation cell forced by surface wave transport, and have little to do with tides.

rotary current *see* § 5.8 and **streams**

satellite altimetry

Remote sensing of the ocean surface height by satellite-mounted microwave radar. Techniques have been developed for extracting the tidal constants for the larger constituents from the satellite data, despite its sampling interval which is generally about twenty times longer than the semi-diurnal period, thus providing an accurate global ocean database of tidal constants, which was previously restricted to areas close to coastal tide gauges and to numerical models which were poorly constrained over wide areas of the ocean surface.

SEAFRAME *see* § 5.1

secondary port

In the context of tide tables, a port for which predictions are required, but for which insufficient data for a reliable harmonic analysis is available and hence, predictions from the nearest standard port (*see* **standard port**) must be used (with suitable corrections). Also called a subordinate port. See also **inference of constituents** (§ 5.9).

secular trend

Long-term trend in any time series, such as one of sea level. "Secular" is usually used to imply a background trend – for example, the trend over several decades of annual sea level – but with the understanding that the secular trend may vary if the length of the time series is significantly extended.

seiche

A standing wave in an enclosed or semi-enclosed body of water set off by weather, seismicity, or incident ocean waves. Seiches are primarily a resonance phenomenon, whose wavelength and period are determined by the geometry. A characteristic feature is the existence of "nodes" - points of minimum water level disturbance, but greatest horizontal currents - and "antinodes", where the reverse occurs. Closed basins (e.g. Lake Geneva) have antinodes at both ends, whereas open basins (e.g., Bay of Fundy) have an antinode at the nearshore end and a node at the open ocean. The Bay of Fundy is frequently cited as an example because it has a natural resonance period close to semidiurnal, and hence the tidal range at the upper end are large due to resonance. The fundamental modes of closed- and open-basin seiches have periods of $2L(gH)^{-1/2}$ and $4L(gH)^{-1/2}$ respectively, where L is the length of the harbour or embayment, H is the mean depth, and g is the gravitational acceleration (9.8 m/s^2). The factor $(gH)^{1/2}$ is the shallow-water wave speed. Higher order seiches of shorter period are also possible, in which case the two formulas must be divided by n and (2n-1) respectively, with n=1,2,3.. in both cases. The formula for the closed-basin mode is known as "Merian's formula".

seismic sea wave *see* tsunami**self-attraction and loading**

The description of the tide-generating potential (§ 1.2) was simplified in several ways, two of which were by ignoring the continuous re-distribution of mass that occurs as the water moves in response to the tidal forces, and by ignoring the warping of the elastic solid earth surface as the water level varies. These effects are commonly combined into the term "self-attraction and loading" (SAL). For each constituent, the global tide is represented as a summation over spherical harmonics. Solutions to the

tidal forcing equations are displacements proportional to the "loading Love numbers" (h'_n , l'_n , and k'_n). The vertical and horizontal displacements are given by h'_n and l'_n respectively, and the effect on the tide-generating potential is given by k'_n . This set of numbers forms the basis for modern computations of SAL. Some early ocean tide models attempted to account for SAL or loading alone by subtracting $\beta\zeta$, where ζ is the water level anomaly, (negative when water level is less than mean), and β typically equalled 0.08 for SAL and 0.03 for loading alone. The advent of accurate global tide models has enabled scientists to establish far more accurate models of SAL which can estimate the appropriate adjustment at each new time step, for each geographic point, for each tidal constituent included in the model (Figure 6.7). *See also Ray (1998) and Baker (1984).*




Figure 6.7 The "SAL tide" for M_2 : (*top*) Amplitude contours in mm; (*bottom*) phase lag (degrees, UT). The phase contours are broadly similar to those seen on maps of the global M_2 tide. Graphic courtesy of Richard Ray. (Note that the SAL function is mathematically defined over land, but is not physically meaningful except over oceans and large lakes.)

semi-diurnal tides *see* **species**

set (of current) *see* **streams**

sidereal day *see* **day**

slack water *see* **streams**

solstitial tides

Tides occurring near the times of summer and winter solstice, when the sun is overhead at the Tropic of Cancer or Capricorn. If the associated constituents (K_1 and P_1) were predominant, the tide would be diurnal, with an annual modulation at their beat frequency, i.e. at a period equal to the tropical year. This being a rarity, the term is of minor currency. These tides seem to have a stronger claim to the term "tropic tide" than those that inherited it, but at least "solstitial" is unambiguous, if tongue-twisting.

species

The Doodson number i_a for each constituent defines its "species". Those of period half-monthly and longer are assigned $i_a = 0$; for periods approximately diurnal, $i_a = 1$; for periods approximately semi-diurnal, $i_a = 2$; for periods approximately ter-diurnal (one-third diurnal), $i_a = 3$; quarter-diurnal, $i_a = 4$; and sixth-diurnal, $i_a = 6$. The different species form distinct groups in a line spectrum.

speed *see* **Section 2.4**

spring high water *see* **tidal planes**

spring low water *see* **tidal planes**

spring tides *see* **fortnightly cycle**

stand of tide *see* **streams**

standard port

In the context of tide tables, a port for which sufficient data is available in order for a set of official predictions to be produced. Also known as a "primary port", especially in North America.

stilling well *see* **tide gauges**


storm surge

The temporary piling-up of water at the coast due to onshore wind and/or low barometric pressure. A storm surge combined with high tide can be particularly dangerous, and even more so in the presence of wind-generated waves. Negative surges (lowered water levels) are also possible. There is a close association between tides and storm surges - the impact of a surge often depends on the state of the tide, and the surge and tide waves may interact over the shelf or as they move up an estuary.

Storm surges are most often caused when a tropical cyclone (also known as a "hurricane" or "typhoon"), generated over the open ocean, moves across the shallower water of the continental shelf. The cyclonic wind circulation is counter-clockwise in the northern hemisphere and clockwise in the southern hemisphere. The strongest onshore winds - and thus, the shoreline with the highest risk of water set-up - is thus to the right (left) of the direction in which the storm is moving, in the northern (southern) hemisphere.

On 2 April 2000, a Tropical Cyclone Tessi struck northeast Queensland, Australia (Figure 6.8), bringing damage to property and uprooting trees. The highest winds, flooding, and greatest damage, was reported at Townsville, to the south of the point of landfall.

Figure 6.8 This map of far north Queensland and the predicted track of TC Tessi was prepared by the Joint Typhoon Warning Center, USNPMOC, Hawaii (reproduced with permission). Times are in "Z" (i.e., UT). The dashed line encircles the 24 hour warning area.



Sea levels and weather data recorded at Cape Ferguson, 20 km south of Townsville, is plotted in Figure 6.9. Fortuitously, the peak gusts and storm surge (sea level residual) nearly coincided with low tide. TC Tessi stalled and veered after the map was prepared - landfall occurred eight hours later, and about 80 km north of the point predicted on the map. Wind direction at the time of maximum speeds was from the southeast (as expected); but by the end of the day it had swung around and was blowing from the northeast. The arrival of the peak gusts after the lowest pressure is a little unusual. It may be due to the cyclone veering and stalling before making landfall, or perhaps the coastal topography.




Figure 6.9
 Weather and water level data recorded by the National Tidal Centre (Bureau of Meteorology) SEAFRAME gauge at Cape Ferguson. Barometric pressure and peak gust are recorded on the hour; sea level is recorded at six minute intervals. Time is UT.

If the alongshore progression of a tropical cyclone is close to the speed of a long wave over the local shelf, resonance may occur and a shelf wave generated. This can dramatically enhance the size of the storm surge. The storm surge is thus a response to a combination of factors - high winds piling up water against the coast, the **inverse barometer effect**, and the resonant shelf wave.

strand line: The high water mark on a shoreline, typified by the presence of flotsam and jetsam.

streams

Same as tidal currents, although some hydrographic authorities use "streams" to refer exclusively to the tidal currents along the principal directions of ebb and flood (which may not differ by 180°, but usually do!). The *set* of the current is the direction in which it flows. On the incoming tide, the streams are said to be in *flood*; the outgoing streams are in *ebb*. The *stand* of the tide occurs near high and low water when the

water level is unchanging. The analogous term for streams is the *slack water* that may or may not occur at the same time. Tidal streams which flow back and forth along a line are *rectilinear*, whereas those that follow an elliptical circuit (due to the coriolis force) are *rotary* (see § 5.8). The ellipse traced out by a tidal current vector in a rotary flow regime is called a *tidal ellipse*. The principal directions of ebb and flow can be computed by the methods of § 5.8.

synodical month *see* **month**

syzygy

The condition whereby the sun, earth and moon are in alignment. *See also* quadrature. The term syzygy is surely used far less often by tidallists than by players of word games.

TASK-2000 *see* **tide software packages**

ter-diurnal *see* **species**

thermocline

A layer in which the temperature decreases significantly (relative to the layers above and below) with depth. The principal ones are designated diurnal, seasonal, and main thermocline. A common feature of thermoclines is the presence of internal tides.

tidal bore

A tidal bore is a moving hydraulic jump caused by tidal propagation up a river. Hydraulic jumps are familiar as the abrupt changes in water level that occur, for example, in a gutter after a heavy rain. In that case, they are stationary, whereas others, such as tidal bores, may propagate as a special type of wave. One of the world's largest tidal bores, on the Qiantang (formerly spelled "Tsien Tang") River near Hangchow, China, has been known to reach nearly ten metres (Figure 6.10).

"Not all bores are boring."

Figure 6.10 Tidal bore on Qiantang River. Photo credit: Dr. J.E. Jones, Proudman Oceanographic Institute, UK.



tidal datum *see* § 5.9

tidal plane *see* § 5.9

tidal prism

Where the tide moves up and down the lower reaches of a river, a volume, known as the tidal prism, of fresh water is displaced each tidal cycle. The tidal prism takes its name from the fact that the front between fresh and salt water is often inclined to the vertical, with the downstream edge of the fresh water riding over the salt.

tidal wave

The response of the ocean to the gravitational forcing of the sun and moon includes the generation of various types of large-scale waves, generically called tidal waves. Sometimes this term is used incorrectly as a synonym for tsunami.

tidal pumping

This term is used in various contexts, including those of coastal aquifers and the bringing of nutrient-rich offshore water into the shallower regions. In the former case, the rise and fall of the tide is often accompanied by a delayed and reduced oscillation of water level in nearby wells. The latter context usually involves a relatively large flood tide bringing water up and into a bay or other semi-enclosed area, where it mixes with water from previous high tides before draining more slowly back to the open ocean. Submarine canyons across the continental shelf may also cause a rectified flow with a net increase of nutrients in the upper layer.

Time and tide: the English word for "time" goes back to an ancient Indo-European form used about six thousand (6000) years ago: "dai-". By the time people were speaking Germanic, about two thousand years ago, "dai-" was being used in two Germanic words: "tídiz" (meaning "a division of time", and "tímon" (meaning something like "an appropriate time [at which to do something]"). The "tídiz" word became Old English "tíd" and then finally "tide".

- Adapted from Word Lore, <http://hea-www.harvard.edu/ECT/Words/>.

time scales *see* § 5.9

tropic tides

At latitudes near the maximum declination of the moon (which varies between 18.3° and 28.6° latitude north and south over the course of the nodal cycle) the diurnal tides are greatest when the moon is near maximum declination. These so-called "tropic tides" are the equivalent of the more common spring tides, with the beat frequencies being diurnal (eg. O_1 and K_1) instead of semi-diurnal. The beat period for O_1 and K_1 is 13.66 days. The tides at Karumba, Queensland, which are dominated by O_1 and K_1 , exhibit this pattern (Figure 6.11). The term may be slightly misleading in that the "Tropics" on the earth are the latitudes $\pm 23.5^\circ$ where the *sun*'s declination reaches its maximum (*see solstitial tides*); nevertheless, the Tropics are also the maximum declinations of the moon when averaged over a nodal cycle. The range (peak to peak distance between high and low tide) at the time when these diurnal tides are greatest is known as the tropic range (*vide*).




Figure 6.11 Predicted tidal heights at Karumba, Queensland (latitude 17° 29' S) over a one month period. Full and new moons are indicated. The beat cycle is caused by O_1 and K_1 .

tropic ranges *see* **tropic tides**

tsunami

Not a tidal term, but included because it is sometimes incorrectly called a “tidal wave”. A tsunami is an ocean wave caused by a disturbance such as an undersea earthquake or landslide, whose wavelength is long compared to the water depth.

twenty-nine day analysis

The analysis of 29 days of tidal observations. 29 days contains a near-integral multiple of the periods of the four major constituents, M_2 (period = 0.518 d), S_2 (0.500 d), O_1 (1.076 d), K_1 (0.997 d). *Viz.*: $29/0.518 = 55.98$, $29/0.500 = 56$, $29/1.076 = 26.95$, $29/0.997 = 29.09$. For this reason time-stepping numerical models driven at the boundary by these four constituents are sometimes run for 29 days (following spin-up). Near-integral multiples also occur at 59, 355, and 738 days.

universal time (UT) *see* § 2.5

upwelling

In the context of tides, upwelling (upwards movement of water) can occur as a result of periodic flow over uneven topography, especially submarine canyons on the continental shelf. Upwelling is more often associated with alongshore winds combined with the coriolis effect, or spatially divergent wind fields over the open ocean, but tidal upwelling can also lead to significant flux of nutrients into the photic zone.

Van de Castele test

A test designed to detect flaws in the mechanical operation of tide gauge chart recorders. A measurement is taken of the positive distance between a fixed point near the top of the gauge, down through the stilling well to water level. The sum of this

distance, which is a maximum at low tide, and the tide gauge reading should be constant through a full tidal cycle. The sum when plotted against the measured distance (with the latter plotted on the vertical axis) should therefore be a vertical line. Deviations from the straight line can be interpreted as faults such as backlash in the gauge mechanism, scaling error, etc. A full description of the test and interpretation is available online from UNESCO/IOC Manual 14:
http://www.pol.ac.uk/psmsl/manuals/ioc_14i.pdf.

vanishing tide *see* **dodge tide**

variational *see* **evection and variation**

year

Four different types of year are of significance to tides. The *sidereal* year is the period taken by earth complete a single orbit of the sun, 365.2564 mean solar days (msd). The *tropical* year, which is measured in relation to the beginnings of the various seasons (specifically, successive vernal equinoxes), is slightly shorter than the sidereal year as a consequence of precession. The axis of the earth is tilted at about $23\frac{1}{2}^{\circ}$ degrees to the perpendicular of the orbital plane. The axis slowly precesses about the perpendicular, in the manner of a “sleeping top”. If it completed a single precession in one day, then we would experience four seasons in a single day. Of course, this is not the case – 26,000 years are required for each precession. This means that the seasons advance $1/26,000^{\text{th}}$ part per sidereal year faster than they would without precession, and the tropical year is therefore only 365.2422 msd. Finally, there is an *anomalous* year, which is the period between successive perihelions. Just as the anomalistic month is slightly longer than a sidereal month, the anomalistic year, 365.2596 msd, is slightly longer than a sidereal year. Of the three types of year, the anomalistic is of greatest importance in tides. The longitude of the sun (λ_h) undergoes a complete cycle in one tropical year. The final type of year, the Julian year, is a rather artificial construct in comparison to the others. It is defined as 365.25 days.